


PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 5 : G01N 31/00, A01N 31/00 A61K 31/715		A1	(11) International Publication Number: WO 90/08315 (43) International Publication Date: 26 July 1990 (26.07.90)
(21) International Application Number: PCT/US90/00121 (22) International Filing Date: 12 January 1990 (12.01.90) (30) Priority data: 297,012 13 January 1989 (13.01.89) US		(74) Agent: MURRAY, Robert, B.; Armstrong, Nikaido, Marmelstein, Kubovcik & Murray, Suite 1000, 1725 K Street, N.W., Washington, DC 20006 (US). (81) Designated States: AT, AT (European patent), AU, BB, BE (European patent), BF (OAPI patent), BG, BI (OAPI patent), BR, CA, CF (OAPI patent), CG (OAPI patent), CH, CH (European patent), CM (OAPI patent), DE, DE (European patent), DK, DK (European patent), ES, ES (European patent), FI, FR (European patent), GA (OAPI patent), GB, GB (European patent), HU, IT (European patent), JP, KP, KR, LK, LU, LU (European patent), MC, MG, ML (OAPI patent), MR (OAPI patent), MW, NL, NL (European patent), NO, RO, SD, SE, SE (European patent), SN (OAPI patent), SU, TD (OAPI patent), TG (OAPI patent), US.	
(60) Parent Application or Grant (63) Related by Continuation US 297,012 (CIP) Filed on 13 January 1989 (13.01.89) (71)(72) Applicants and Inventors: PANG, Peter, K., T. [US/CA]; 52225 Range Road, 205 Carriage Lane, Sherwood Park, Alberta T8A 2A6 (CA). WANG, Lawrence, C., H. [CA/CA]; 5012-144 St., Edmonton, Alberta T6G 2E9 (CA). BENISHIN, Christina, G. [CA/CA]; 218-53431 Range Rd., 221, Ardrossan, Alberta T0B 0E0 (CA). LIU, Hsing, J. [CA/CA]; 3543-105B St., Edmonton, Alberta T6J 2K9 (CA).		Published With international search report.	

(54) Title: COMPOSITION AND METHOD FOR TREATMENT OF SENILE DEMENTIA



(57) Abstract

Ginsenosides Rb<sub>1</sub> and Rg<sub>1</sub> enhance the availability of acetylcholine in the cortical and hippocampal regions of the brain and alleviate the symptoms of Alzheimer-type senile demential. The Rb<sub>1</sub> or Rg<sub>1</sub> may be administered together with a metabolic precursor for acetylcholine and/or with a cholinesterase inhibitor. Pure Rb<sub>1</sub> is located from a mixture of ginsenosides by a process involving vacuum chromatography on silica gel. Preferably, the mixture of ginsenosides is enriched in Rb<sub>1</sub> by partition between an aqueous system and water ethyl acetate-butanol.

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	ES	Spain	MG	Madagascar
AU	Australia	FI	Finland	ML	Mali
BB	Barbados	FR	France	MR	Mauritania
BE	Belgium	GA	Gabon	MW	Malawi
BF	Burkina Faso	GB	United Kingdom	NL	Netherlands
BG	Bulgaria	HU	Hungary	NO	Norway
BJ	Benin	IT	Italy	RO	Romania
BR	Brazil	JP	Japan	SD	Sudan
CA	Canada	KP	Democratic People's Republic of Korea	SE	Sweden
CF	Central African Republic	KR	Republic of Korea	SN	Senegal
CG	Congo	LI	Liechtenstein	SU	Soviet Union
CH	Switzerland	LK	Sri Lanka	TD	Chad
CM	Cameroon	LU	Luxembourg	TG	Togo
DE	Germany, Federal Republic of	MC	Monaco	US	United States of America
DK	Denmark				

-1-

## COMPOSITION AND METHOD FOR TREATMENT OF SENILE DEMENTIA

This application is a continuation-in-part of application Serial No. 07/297,012, filed January 13, 1989.

5      The present invention is directed to compositions and methods for alleviating the symptoms of senile dementia of the Alzheimer's type. In one specific aspect, the present invention is directed to an improved process for the isolation and purification of a ginsenoside useful in practicing that method.

Background of the Invention

10      Senile dementia of the Alzheimer's type (SDAT) is widely recognized as a problem of increasing proportions in North America as well as around the world. The disease is associated with progressive physical and mental impairment to the point where the patient requires total care, and becomes a tremendous social and  
15      economic burden. Progress of the disease is believed to be related to degeneration of certain nerve tracts in the central nervous system, resulting in the loss of associated functions. Pathological studies indicate that brains of SDAT patients have loss of several neurotransmitter systems, related to different  
20      functions, but the system which is implicated the most is the cholinergic system. Studies show that several important cholinergic tracts innervating the cortical and hippocampal regions degenerate. Although this particular degeneration may not account for all of the symptoms of SDAT, it may account for the  
25      cognitive and memory deficits, which are some of the most difficult symptoms for patients and their families to deal with.

The pharmacological approaches which have been proposed for the managements of SDAT symptoms may be classified in two ways. The first is drugs which improve the function of existing neurons, especially to increase cholinergic nerve function. The second is drugs which decrease degeneration/increase regeneration of nerves.

Two types of drugs have been used in clinical trials to improve central cholinergic functions. The first is compounds which increase the availability of the existing endogenous neurotransmitter, acetylcholine (ACh); and the second is compounds which are exogenous, and mimic the effects of the endogenous transmitter at the receptor. However, these compounds exhibit side effects which limit their use.

It is generally believed that compounds which will increase the availability of the endogenous neurotransmitter are more desirable. Substances in this category are cholinesterase inhibitors, such as physostigmine and pyridostigmine, which decrease the breakdown of ACh, thus prolonging its functional lifetime at the crucial location, the synaptic cleft, and choline and lecithin, which increase the availability of the precursor for synthesis. Thus far, other compounds have not been described which directly increase the availability of the endogenous neurotransmitter ACh by any other mechanisms, except by blockade of inhibitory presynaptic receptors (with e.g. atropine or clonidine), or by non-specific depolarization of nerves (e.g. veratridine).

Ginseng is the name given to the dried roots of the ginseng plants (genus Panax) and, more particularly, to extracts

-3-

of those roots. The roots and their extracts contain a variety of substances including saponins and sapogenins.

Ginseng has been extensively used, mostly in Asia, as a tonic to promote health and well being, and as a medicine in the treatment of various disease conditions. The beneficial attributes of ginseng are attributed to its saponin content, a mixture of glucosides referred to collectively as ginsenosides.

#### Discussion of the Prior Art

U.S. Patent 4,157,894 to Bombardelli discloses the isolation of saponins from ginseng roots and the use of a purified concentrate in the geriatric field for elderly patients having difficulty in digesting less concentrated preparations. Bombardelli also discloses the structures of saponins Rb<sub>1</sub>, Rb<sub>2</sub>, Rc, Rd, Re, Rf and Rg.


U.S. Patent 4,702,949 to Liu discloses a composition comprising 5-15% ginsenoside, 30-50% tetramethylpyrazine, 30-50% astragalan, and 5-15% atractylol in the treatment of cerebral vascular insufficiency and resultant paraplegia, hemiplegia and impaired neurofunction.

U.S. Patents 4,157,894; 4,317,816; 4,446,130; 4,647,460; 4,684,628; 4,687,761; and 4,755,504 disclose the use of ginseng or ginseng extracts, alone, or in combination with other substances for various medically related purposes.

A procedure for the isolation of a crude mixture of ginsenosides for ginseng is described by J. Shoji, "Advances in Chinese Medicinal Materials Research", World Scientific Publishing Company, Singapore, pages 455-469 (1985). Material prepared by this established procedure is commercially available.

The preparation of ginseng extracts is also disclosed by Bombardelli and by Liu, discussed above, and in the other U.S. patents listed above.

Rb<sub>1</sub> and Rg<sub>1</sub> have the structural formula:



In Rb<sub>1</sub>, R<sub>1</sub> is D-glucose B(1→6)D-glucose, R<sub>2</sub> is D-glucose B(1→2)D-glucose and R<sub>3</sub> is H. In Rg<sub>1</sub>, R<sub>1</sub> is D-glucose, R<sub>2</sub> is H, and R<sub>3</sub> is O-D-glucose.

Existing procedures for the isolation and purification of ginsenoside Rb<sub>1</sub> include standard column chromatography, thin-layer chromatography, and high performance chromatography. These methods are laborious for the isolation of the compound in large quantities and often yield a low purity product.

#### Brief Description of the Invention

A primary object of the present invention is to provide compositions and methods for the treatment of senile dementia of the Alzheimer's type.

Another object of the invention is to provide an improved process for the isolation and purification of a ginsenoside used in that method for treating senile dementia.

We have discovered that ginsenosides Rb<sub>1</sub> and Rg<sub>1</sub> directly and selectively increase acetylcholine function in the brain and,

accordingly, are useful in alleviating the symptoms of senile dementia. Those ginsenosides may be administered together with metabolic precursors for ACh synthesis and/or with cholinesterase inhibitors.

5

In one specific aspect, the present invention is a method for alleviating the symptoms of Alzheimer-type senile dementia, ~~which comprises administering to a mammal affected with Alzheimer-~~ type senile dementia an amount of ginsenoside Rb<sub>1</sub> or of ginsenoside Rg<sub>1</sub> effective to increase the availability of acetylcholine in the cortical and hippocampal regions in the brain of the mammal.

10

In a second specific aspect, the present invention is a process for the isolation of ginsenoside Rb<sub>1</sub> which comprises the steps:

15

(a) dissolving a mixture of ginsenosides in methyl alcohol;

20

(b) adsorbing the mixture of ginsenosides on silica gel by containing the solution of the mixture in methyl alcohol with the silica gel and evaporating the alcohol;

25

(c) placing the silica gel having the mixture of ginsenosides adsorbed thereon in a column for vacuum chromatography prepacked with silica gel;

(d) passing a mixture of chloroform and methyl alcohol through the columns to elute ginsenoside Rb<sub>1</sub>; and

(e) recovering ginsenoside Rb<sub>1</sub> from the chloroform and methyl alcohol eluate.

-6-

A mixture of ginsenosides enriched in Rb<sub>1</sub> particularly useful as the starting material in the process of the present invention may be obtained by:

- 5 (a) dissolving a mixture of crude ginsenosides obtained by extraction of ginseng in water;
- (b) washing the aqueous solution of mixed ginsenosides with ethyl acetate;
- 10 (c) extracting the ethyl acetate-washed solution successively with a mixture of 4 volumes of ethyl acetate and 1 volume of 1-butanol, with 1 volume of ethyl acetate and 1 volume of 1-butanol, and with 1-butanol pre-saturated with water;
- (d) combining the ethyl acetate-1-butanol and 1-butanol-water extracts; and
- 15 (e) recovering a mixture of ginsenosides enriched in Rb<sub>1</sub> from the combined extracts.

The vacuum chromatographic process described above yields ginsenoside Rb<sub>1</sub> of good quality and can be adapted to the preparation of larger amounts of that ginsenoside. Exceptionally  
20 high purity Rb<sub>1</sub> can be obtained by using a mixture of ginsenosides enriched in Rb<sub>1</sub> as the starting material in the process.

In a third specific aspect, the preset invention is a composition for alleviating the symptoms of Alzheimer-type senile dementia comprising:

- 25 a) 25-250 mg of Rb<sub>1</sub> or Rg<sub>2</sub> together with a pharmaceutically acceptable carrier therefore,
- b) 25-250 mg of Rb<sub>1</sub> or Rg<sub>2</sub> and a metabolic precursor for acetylcholine, or



-7-

- c) 25-250 mg of  $Rb_1$  or  $Rg_2$  and a cholinesterase inhibitor.

#### Brief Description of the Drawings

Figure 1 is a graph of acetylcholine release plotted against log concentration  $Rb_1$  at high and low concentrations of potassium.

Figures 2 and 3 are graphs plotting electrically stimulated  $^3H$ -acetylcholine release against time.

Figures 4 and 5 are a graphs of choline uptake plotted against log concentration  $Rb_1$  and against log concentration  $Rg_1$ , respectively.

Figure 6 is a graph of  $^{45}Ca$  uptake at high and low concentrations of potassium.

Figure 7 is a graph showing the effects of  $Rb_1$  on calcium uptake at nerve endings.

Figure 8 is a graph showing the effects of  $Rb_1$  on intracellular calcium concentration in cultured neuroblastoma cells.

Figure 9 is a graph illustrating that  $Rb_1$  itself has no anticholinesterase (AChE) activity.

Figure 10 is a graph showing the inability of  $Rb_1$  to displace QNB from nerve ending receptors.

Figures 11 and 12 are graphs illustrating that the increase in ACh content at nerve endings is not caused by an increase in activity choline acetyltransferase (ChAT).

Figure 13 is a graph illustrating that the increase in the release of ACh is accompanied by an increase in the uptake of

-8-

choline into the nerve endings. Figures 14 and 15 are graphs illustrating the kinetics of that uptake.

Figure 16 is a graph illustrating that  $Rb_1$  does not displace radio labelled HC-3 and did not increase the number  
5 binding sites in the brain.

Figure 17 is a graph illustrating that the administration of  $Rb_1$ , increases the number of choline-uptake sites in the rat.

#### Detailed Description of the Invention

Our invention is further illustrated by means of the  
10 following non-limiting examples:

##### Example 1: (isolation of $Rb_1$ )

About 1 g of crude ginsenosides obtained by the procedure described by Shoji in "Advances in Chinese Medicinal Materials Research" was dissolved in 10 ml of methanol. The resultant  
15 solution was mixed with 10 g of silica gel (Merck 0.040-0.063 mm particle size, 230-400 mesh ASTM). The silica gel was air dried for about 1 hour and then placed in a column for vacuum chromatography, as described by Coll et al, Aust. J. Chem., 30, 1305 (1977), prepacked with 80 g of silica gel. Elution with a  
20 solution of chloroform and methanol (85:15) gave a fraction from which 75mg of > 98% pure  $Rb_1$  was obtained. An additional 120 mg of  $Rb_1$ , purity about 60%, was recovered from other fractions of the eluate.

##### Example 2: (preparation of ginsenosides enriched in $Rb_1$ )

25 600 mg of the crude mixture of ginsenosides obtained by the procedure of Shoji was dissolved in 10 ml water. The aqueous solution was washed with ethyl acetate (2 x 40 ml) and then extracted with a solution of ethyl acetate and 1-butanol (4:1; 4

x 40 ml) followed by extraction with a 1:1 solution of ethyl acetate-1-butanol (2 x 40 ml) and then 1-butanol presaturated with water (2 x 40 ml). The last three extracts were combined and concentrated to give 140 mg of ginsenosides containing about 60% of Rb<sub>1</sub>. Use of that enriched mixture as the starting material in Example 1 gave essentially pure Rb<sub>1</sub>.

The following examples illustrate that Rb<sub>1</sub> and Rg<sub>1</sub> directly and selectively increase acetylcholine function in the brain and are useful in the treatment of Alzheimer-type senile dementia.

10      Example 3:

Pinched off nerve endings (synaptosomes) from whole rat brain were first incubated in the presence of the precursor <sup>3</sup>H-choline, which is converted intracellularly to <sup>3</sup>H-ACh. The release of ACh from synaptosomes was quantitated under low [K] and high [K] conditions intended to simulate physiological stimulation. The addition of Rb<sub>1</sub> increased the release of <sup>3</sup>H-ACh as shown in Fig. 1.

Example 4:

Using a different protocol, synaptosomes, which had been previously incubated with <sup>3</sup>H-choline, as in the first series of experiments, were continually perfused with HEPES-buffered Krebs solution. The release of ACh was effected by subjecting the synaptosomes to electrical field stimulation, which more closely mimics physiological stimulation. In Fig. 2 and 3, the left panel shows the release pattern in the absence of any added drug, and the right panel shows the release pattern in the presence of 10<sup>-6</sup>M drug during the last 25 minutes of perfusion. The results obtained by calculating the ratio of the amounts of ACh released during the two periods of electrical stimulation (the areas under

the curves, S2/S1) indicate that both Rb<sub>1</sub> and Rg<sub>1</sub> increase the electrically stimulated release. In addition, Rg<sub>1</sub> stimulates the resting release of <sup>3</sup>H-ACh, as indicated, by the increase in the resting afflux of <sup>3</sup>H-ACh when the drug was first added. While the net amount of <sup>3</sup>H-ACh released in each run can vary, depending on the amount of protein on the filter and aging of the synaptosomes, the S2/S1 ratio remains quite constant from run to run using a given chamber. Therefore, in each experiment the same chamber was used for both control and test-drug runs.

The stimulation of ACh release is also associated with an increase in the specific uptake of the precursor <sup>3</sup>H choline as shown in Figs. 4 and 5. Although the magnitude of stimulation of choline uptake is not as great as the magnitude of stimulation of ACh release, it is a consistent and significant effect, and can still quantitatively account for the increase in release. This suggests a general stimulation of brain cholinergic function. Further, preliminary experiments show that of three cholinergic brain regions examined, cortex, stratum, and hippocampus, stimulation of ACh release from synaptosomes is most pronounced in the hippocampus, a brain region strongly implicated in memory functions, and to a lesser extent in the cortex.

Example 5:

This example confirms that Rb<sub>1</sub> and Rg<sub>1</sub> do not stimulate the release of neurotransmitter by non-selectively depolarizing nerve endings. The effects of Rb<sub>1</sub> on the resting voltage dependent uptake of <sup>45</sup>Ca into rat brain synaptosomes are shown in Fig. 6. these results indicate that the dramatic stimulation of <sup>3</sup>H-ACh release is associated with only a minimal increase in the resting uptake of Ca, upon which ACh release is dependent. If this

-11-

compound were acting as a non-specific depolarizing agent, one would predict the resting uptake of  $^{45}\text{Ca}$  would be stimulated several fold, to the same level as the depolarized (52.5 K) uptake.

5        Example 6:

Table 1 below contains data illustrating the effect of  $\text{Rb}_1$  on the release of acetylcholine from hippocampal tissue, a brain region which is associated with memory and learning. These data indicate that  $\text{Rb}_1$  can stimulate the release of ACh from brain tissue. This effect is observed in the presence or absence (10 mM EGTA) of calcium, which may suggest that the source of the stimulated release of ACh is not from the vesicular pool, as is normally the case, but from the cytoplasmic pool. Furthermore the stimulation of release in the absence of calcium is more pronounced in the simultaneous presence of a cholinesterase inhibitor, paraxon. This type of drug is also known to increase the amount of ACh in the cytoplasm, by virtue of its ability to inhibit intracellular AChE, and this observation would be consistent with the notion that the source of ACh which is stimulated by  $\text{Rb}_1$  is the cytoplasm.

Table 1

CONDITION	RATIO	N
25        with Paraaxon		
Control	1.053±0.058	29
10 mM EGTA	0.299±0.062	10
10 <sup>-7</sup> M $\text{Rb}_1$	2.107±0.284	14
10 <sup>-6</sup> M $\text{Rb}_1$	1.380±0.112	10
10 <sup>-6</sup> M $\text{Rb}_1$ +		
10 mM EGTA	1.139±0.338	9

-12-

without Paraoxon

5	Control	1.022±0.024	24
	10 mM EGTA	0.357±0.126	6
	10 <sup>-7</sup> M Rb <sub>1</sub>	2.108±0.542	11
	10 <sup>-6</sup> M Rb <sub>1</sub>	1.428±0.169	10
	10 <sup>-6</sup> M Rb <sub>1</sub> + 10 mM EGTA	0.676±0.194	9

Example 7:

The stimulation of ACh release is not accompanied by an increase in intracellular calcium in nerves. Figure 7 shows the effects of Rb<sub>1</sub> on nerve ending calcium uptake, and Figure 8 shows the effects of Rb<sub>1</sub> on the intracellular calcium concentration in cultured neuroblastoma cells.

Example 8:

Figure 9 shows that Rb<sub>1</sub> itself also has no anticholinesterase activity.

Example 9:

Table 2 below summarizes the effect of Rb<sub>1</sub> on the intracellular stores of choline and ACh when tissues are incubated in low or high potassium, and in the absence or presence of Rb<sub>1</sub>. Rb<sub>1</sub> produces an increase in and in the absence or presence of Rb<sub>1</sub>. Rb<sub>1</sub> produces an increase in the total <sup>3</sup>H contents (choline and ACh) and in the ACh contents of the cytoplasmic fraction (S3), but not in the vesicular (P3) fraction.

Table 2Total <sup>3</sup>H contents of Subcellular Fractions

	<u>S3</u>	<u>P3</u>
Low K	166.3±20.1	55.1±4.2
Low K + Rb <sub>1</sub>	214.4±18.9	62.0±5.5
High K	209.0±19.5	87.3±7.3
High K + Rb <sub>1</sub>	216.0±21.8	97.2±8.8

<sup>3</sup>H ACh Contents of Subcellular Fractions

	<u>S3</u>	<u>P3</u>
Low K	38.6±9.4	7.0±0.72
Low K + Rb <sub>1</sub>	48.5±6.9	7.8±0.3
High K	16.8±0.9	8.0±0.9
High K + Rb <sub>1</sub>	17.8±0.9	6.8±0.7

Example 10:

The stimulation in activity is not mediated by autoreceptors. Figure 10 shows the lack of ability of Rb<sub>1</sub> to bind to nerve ending muscarinic receptors (displace QNB from its nerve ending receptor).

Example 11:

The increase in the nerve ending ACh content is not caused by an increase in the activity of the synthetic enzyme choline acetyltransferase (ChAT) as is shown in Figures 11 and 12.

Example 12:

The increase in the release of ACh is accompanied by an increase in the uptake of the precursor choline into nerve endings, which is shown in Figure 13. The kinetics of the increase in uptake are presented in Figures 14 and 15, and the kinetic contents are presented in Table 3. These results indicate that the increase in choline uptake is not due to an increase in the affinity of the carrier for the substrate choline, but to an increase in the maximum velocity of the carrier.

Table 3

	<u>K<sub>m</sub></u>	<u>V<sub>max</sub></u>
Control	12.74 $\mu$ M	475
10 <sup>-10</sup> Rb <sub>1</sub>	6.037 $\mu$ M	709.4
10 <sup>-8</sup> Rb <sub>1</sub>	31.1 $\mu$ M	4572

The mechanism of the increase in the velocity of the carrier could be accounted for in two ways (1) an increase in the turnover rate of the carriers, or (2) in increase in the number of carriers in the plasma membrane. In order to discriminate between these two possibilities, the number of carrier sites was probed with radiolabelled hemicholinium-3 (HC-3). In in vitro administration experiments, presented in Figure 16, Rb<sub>1</sub> did not displace radiolabelled HC-3 (unlabelled HC-3 was also used as a displacer, as a positive control), and did not increase the apparent number of binding sites. Initially, Rb<sub>1</sub> seems to increase the uptake of choline by simply increasing the turnover rate of the carriers. When rats were administered with Rb<sub>1</sub> (5mg/kg/day) for three days, there was an increase in the number of choline uptake sites, suggesting that chronic administration of the compound increases the number of carriers (Figure 17).

When using Rb<sub>1</sub> and Rg<sub>1</sub> to alleviate the symptoms of Alzheimer-type senile dementia, Rb<sub>1</sub> and Rg<sub>1</sub> can be processed by conventional methods of galenic pharmacy into pharmaceutical preparations for oral or parenteral administration, e.g., to mammals including humans. Conventional excipients are pharmaceutically acceptable organic or inorganic carrier substances suitable for parenteral, enteral or topical application which do not deleteriously react with Rb<sub>1</sub> or Rg<sub>1</sub>. Suitable pharmaceutically acceptable carriers include but are not limited to water, salt solutions, alcohols, gum arabic, vegetable oils, polyethylene glycols, gelatine, lactose, amylose, magnesium stearate, talc, silicic acid, viscous paraffin, perfume oil, fatty acid monoglycerides and diglycerides, pentaerythritol fatty acid esters, hydroxy-methylcellulose, polyvinyl pyrrolidone, etc. The



-15-

pharmaceutical preparations can be sterilized and if desired mixed with auxiliary agents, e.g., lubricants, preservatives, stabilizers, wetting agents, emulsifiers, salts for influencing osmotic pressure, buffers, coloring, flavoring and/or aromatic substances and the like which do not deleteriously react with Rb<sub>1</sub> and Rg<sub>1</sub>.

For parental application, particularly suitable are injectable sterile solutions, preferably oily or aqueous solutions, as well as suspensions, emulsions, or implants, including suppositories.

For enteral application, particularly suitable are tablets, dragees, suppositories or capsules having talc and/or a carbohydrate carrier or binder or the like, the carrier preferably being lactose and/or corn starch and/or potato starch.

A syrup, elixir or the like can be used wherein a sweetened vehicle is employed. Sustained release compositions can be formulated including those wherein Rb<sub>1</sub> and Rg<sub>1</sub> is protected with differential degradable coatings, e.g., by microencapsulation, multiple coatings, etc.

Depending on the type of mammal to which it is being administered, the daily dosage of Rb<sub>1</sub> or Rg<sub>1</sub> for a mammal weighing 50 kg is generally about 100 - 1000 mg. preferably administered 3 or 4 times a day in divided doses. Thus, a suitable dosage form would contain 25 - 250 mg of Rb<sub>1</sub> or Rg<sub>1</sub>. Suitable daily dosages of metabolic precursors of acetylcholine, and of cholinesterase inhibitors, are well known to those skilled in the art. For example, acetylcholine precursor, such as choline, usually as its chloride or bitartrate, dimethylaminoethanol, a synthetic precursor of choline, phosphatidylcholine and lecithin, are

-16-

generally administered in amounts ranging from 5 to 50 g/day. Choline and lecithin are sometimes administered in conjunction with nootropic agents such as piracetam, analogues of piracetam and aminopyridines. As for acetylcholinesterase inhibitors, the  
5 usual dose for neostigmine is 15-30 mg, for pyridostigmine 60-180 mg, for ambenonium 10-20 mg, and for tetrahydroacridine 25-150 mg.

When Rb<sub>1</sub> or Rg<sub>1</sub> is administered together with a metabolic precursor acetylcholine and/or a cholinesterase inhibitor, the dosage form  
10 will generally contain the usual dosage of the acetylcholine precursor and/or the usual dosage of the cholinesterase inhibitor. Appropriate dosages and regimens for a given host can be determined using conventional considerations, e.g., by customary comparison of the differential activities of the subject compound  
and of a known agent, e.g., by means of conventional  
15 pharmacological protocols.

WHAT IS CLAIMED IS:

1. A process for the isolation of ginsenoside  $Rb_1$ , which comprises the steps:

(a) dissolving a mixture of ginsenosides in methyl alcohol;

(b) adsorbing the mixture of ginsenosides on silica gel by contacting the solution of the mixture in methyl alcohol with the silica gel and evaporating the alcohol;

(c) placing the silica gel having the mixture of ginsenosides adsorbed thereon in a column for vacuum chromatography prepacked with a fresh portion of silica gel;

(d) passing a mixture of chloroform and methyl alcohol through the column to elute ginsenoside  $Rb_1$ ; and

(e) recovering ginsenoside  $Rb_1$  from the chloroform and methyl alcohol eluate.

2. A process for the preparation of a mixture of ginsenosides enriched in  $Rb_1$  which comprises the steps:

(a) dissolving a crude mixture of ginsenosides obtained by extraction of ginseng in water;

(b) washing the aqueous solution of mixed ginsenosides with ethyl acetate;

(c) extracting the ethyl alcohol-washed aqueous solution successively with a mixture of 4 volumes of ethyl acetate and 1 volume of 1-butanol, with an equivolume mixture of ethyl acetate and 1-butanol, and with 1-butanol pre-saturated with water;

(d) combining the ethyl acetate-1-butanol and 1-butanol-water extracts; and

(e) recovering a mixture of ginsenosides enriched in Rb<sub>1</sub> from the combined extracts.

3. A process for the preparation of purified Rb<sub>1</sub> which comprises using the ginsenoside mixture enriched in Rb<sub>1</sub> obtained by the process of claim 2 as the mixture of ginsenoside in the process of claim 1.

4. A method for alleviating the symptoms of Alzheimer-type senile dementia, which comprises administering to a mammal affected with Alzheimer-type senile dementia an amount of ginsenoside Rb<sub>1</sub> or of ginsenoside Rg<sub>1</sub> effective to increase the availability of acetylcholine in the cortical and hippocampal regions in the brain of the mammal.

5. A method according to claim 5, wherein the Rb<sub>1</sub> or Rg<sub>1</sub> is administered to the mammal in a daily dosage of 100 - 1000 mg.

6. A method according to claim 6, wherein the daily dosage is administered in portions 3 or 4 times per day.

7. A method according to claim 4, wherein the Rb<sub>1</sub> or Rg<sub>1</sub> is administered together with a metabolic precursor for acetylcholine.

8. A method according to claim 7, wherein the metabolic precursor is lecithin or choline.

9. A method according to claim 4, wherein the Rb<sub>1</sub> or Rg<sub>1</sub> is administered together with a cholinesterase inhibitor.

10. A method according to claim 9, wherein the inhibitor is physostigmine, pyridostigmine or paraoxon.

---

11. A method according to claim 4, wherein the Rb<sub>1</sub> or Rg<sub>1</sub> is administered together with a metabolic precursor for acetylcholine and a acetylcholinesterase inhibitor.

12. A composition for alleviating the symptoms of Alzheimer-type senile dementia comprising 25-100 mg of Rb<sub>1</sub> or Rb<sub>2</sub>.

13. A composition according to claim 12 additionally comprising a metabolic precursor for acetylcholine.

14. A composition according to claim 12 additionally comprising a cholinesterase inhibitor.

15. A composition according to claim 13 additionally comprising a cholinesterase inhibitor.




FIG. 1

SUBSTITUTE SHEET




FIG. 2B




FIG. 2A

3 / 15




FIG. 3A




FIG. 3B

SUBSTITUTE SHEET






FIG.6




FIG.5




FIG.4

SUBSTITUTE SHEET

5 / 15




FIG. 7

SUBSTITUTE SHEET

6 / 15




FIG. 8

SUBSTITUTE SHEET

7/15




FIG. 9

8 / 15





FIG. 10

SUBSTITUTE SHEET

9 / 15



-LOG ( Rb1 )

FIG. II

SUBSTITUTE SHEET

10 / 15




FIG. 12

SUBSTITUTE SHEET

11 / 15


 $\text{LOG}(1/100)$ 

FIG. 13

SUBSTITUTE SHEET






FIG. 14

SUBSTITUTE SHEET

13 / 15




FIG. 15

SUBSTITUTE SHEET

14 / 15




FIG. 16

SUBSTITUTE SHEET

15 / 15




FIG. 17

SUBSTITUTE SHEET

# INTERNATIONAL SEARCH REPORT

International Application No. PCT/US90/00121

<b>I. CLASSIFICATION OF SUBJECT MATTER</b> (If several classification symbols apply, indicate all) <sup>6</sup>		
According to International Patent Classification (IPC) or to both National Classification and IPC U.S. 23/230R 514/54&879 IPC. (5) G01N 31/00, A01N 31/00 CL. 536/5, 127&128 424/195.1 A61K 31/715		
<b>II. FIELDS SEARCHED</b>		
Minimum Documentation Searched <sup>7</sup>		
Classification System	Classification Symbols	
U.S.	23/230R 536/5, 127&128 424/195.1 514/54&879	
Documentation Searched other than Minimum Documentation to the extent that such documents are included in the fields searched <sup>8</sup>		
<b>III. DOCUMENTS CONSIDERED TO BE RELEVANT <sup>9</sup></b>		
Category <sup>*</sup>	Citation of Document, <sup>11</sup> with indication, where appropriate, of the relevant passages <sup>12</sup>	Relevant to Claim No. <sup>13</sup>
A	US, A, 4,157,894 (BOMBARDELLI) 12 June 1979 (See entire document)	1-15
A	US, A, 4,317,816 (ARICHI ET AL) 02 MARCH 1982 (See entire document)	1-15
X A	US, A, 4,339,442 (TAKEMOTO ET AL) 13 July 1982 (See entire document)	1-3 4-15
A	US, A, 4,446,130 (HACHIYA ET AL) 01 MAY 1984. (See entire document)	1-15
X A	US, A, 4,621,137 (MIYAKE ET AL) 04 November 1986. (See entire document)	12 1-11 and 13-15
A	US, A, 4,647,460 (LEE) 03 March 1987. (See entire document)	1-15
A	US, A, 4,684,628 (LIU) 04 August 1987. (See entire document)	1-15
<p><sup>*</sup> Special categories of cited documents: <sup>10</sup></p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"&amp;" document member of the same patent family</p>		
<b>IV. CERTIFICATION</b>		
Date of the Actual Completion of the International Search	Date of Mailing of this International Search Report	
12 MARCH 1990	18 APR 1990	
International Searching Authority	Signature of Authorized Officer	
ISA/US	RONALD W. GRIFFIN	

III. DOCUMENTS CONSIDERED TO BE RELEVANT (CONTINUED FROM THE SECOND SHEET)		
Category *	Citation of Document, with indication, where appropriate, of the relevant passages	Relevant to Claim No
A	US, A, 4, 687,761 (LIU) 18 August 1987 (See entire document)	1-15
<u>X</u> A	US, A, 4,755,504 (LIU) 05 July 1988 (See entire document)	<u>12-</u> 1-11 and 13-15
A,P	US, A, 4,814,339 (ROTONDO) 21 March 1989 (See entire document)	1-15
A,P	US, A, 4,837,219 (HUTTERER) 06 June 1989 (See entire document)	1-15
A, P	US, A, 4,847,082 (SABIN) 11 July 1989 (See entire document)	1-15
A,P	US, A, 4,851,414 (SHIOZAKI ET AL) 25 July 1989 (See entire document)	1-15